

University of Groningen

Long-term effects of large and small herbivores on plant diversity in a salt-marsh system

Chen, Qingqing

DOI:
[10.33612/diss.111645595](https://doi.org/10.33612/diss.111645595)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Chen, Q. (2020). *Long-term effects of large and small herbivores on plant diversity in a salt-marsh system*. [Thesis fully internal (DIV), University of Groningen]. <https://doi.org/10.33612/diss.111645595>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Propositions

Accompanying the dissertation

Long-term effects of large and small herbivores on plant diversity in a salt-marsh system

Qingqing Chen

1. Long-term management is needed for conserving plant diversity in some naturally developed salt marshes (Chapter 2).
2. Low-moderate densities (usually relative to the productivity of the site) of large domestic herbivores, or wild ones, would play a positive role in conserving plant diversity in the long term in grasslands (Chapter 3).
3. Long-term non-trophic effects of large herbivores on plant diversity are underestimated. Non-trophic effects (e.g. trampling, deposition of urine and dung) of large herbivores increased over time, exceeding that of trophic effects (aboveground biomass consumption) 23 years after the start of the experiment in a salt marsh (Chapter 3).
4. Livestock (e.g. dairy cows) are increasingly being kept indoors (Mandel *et al.* 2016), fed by mown grasses and crops. Grasslands are mowed or converted to croplands, in which non-trophic effects of large herbivores have been removed, which has, at least, contributed to the decline in plant diversity (O'Mara 2012; Tschamntke *et al.* 2012; Chapter 3).
5. Small herbivores can have a strong and long-lasting impact on plant diversity, but it highly depends on their abundance, which in turn depends on the quality and abundance of forage plants (Chapter 4).
6. Grazing may lead to increased homogeneity in genetic population structure at the landscape scale (Smith *et al.* 2009; Chapter 5).
7. People should not only complain what they want to but cannot have, but also fight for it.
8. It is not the most intellectual of the species that survives; it is not the strongest that survives; but the species that survives is the one that is able best to adapt and adjust to the changing environment in which it finds itself (Leon C. Megginson).